


Bloating After Eating

Bloating is often linked to overeating and heavy meals. No doubt, the quantity and option of food is a significant factor that can contribute to bloating, however did you understand that it can also be brought on by certain medical conditions? Even some lifestyle factors are connected to bloating and stomach discomfort after consuming. Periodic bloating might not constantly show an underlying illness. Nevertheless, if it occurs regularly or if the signs are severe, you should seek medical interest.



Causes of Bloating - Medical Conditions

Food intolerance: Bloating is one of the classic signs of food intolerance, which may also cause nausea, abdominal pain, [diarrhea](#) or [constipation](#). This condition must not be puzzled with food allergy. Food intolerance symptoms generally appear within a couple of hours or days after consumption. This condition is triggered by the body's inability to produce the specific enzyme required for digesting that food. It is primarily connected with foods, like wheat (gluten intolerance), dairy products, soy products, shellfish, eggs, peanuts and chocolate.

Peptic Ulcers

Repeating episodes of stomach pain or discomfort might be because of the swelling of the stomach lining. This lining performs the function of protecting the stomach from the acids. Long term use of specific anti-inflammatory medications, extreme consumption of alcohol and smoking could also cause erosion in stomach lining. Due to swelling and erosion of the thick mucous layer or the stomach lining, hydrochloric acid and pepsin present in the stomach can trigger sores or peptic ulcers. Ulcers might also be triggered due to an infection in the stomach lining by a bacterium referred to as H. pylori.

- Whenever you are feeling things, which you believe is not regular, then seeking advice from a physician ought to constantly be on top of your list.
- Do not make excuses on not going.
- If your body is offering you indications of illness then understanding what should be done is very important.
- Know what you are experiencing, exactly what triggers it and the best ways to cure it will definitely provide you a better life ahead of you.

Why Do You Feel Bloating

Bloating is frequently related to tightness in the stomach, flatulence and burping. Mostly, this happens due to the existence of big quantities of gas in the stomach. Typically, a small amount of gas is produced in the intestinal tracts, as a by-product of digestion. In many cases, a big quantity of gas is produced in the digestive tract, due to different causes like, disability of the digestion process. This increases the pressure inside the abdomen, which may broaden to accommodate the contents.

This Describes the Inflated Belly, Due to Bloating

The boost in pressure causes the tight feeling and the periodic cramping and discomfort. Stomach pain caused by bloating may radiate to any other place, like the chest and back. The body attempts to expel the gas through belching and by passing wind. The signs may go away in a few hours or take a couple of days, according to the seriousness of the condition and the underlying cause. Some people have the tendency to swallow excess air into the stomach, resulting in bloating. Even physical blockages (like tumors) in the intestines might cause bloating. The causes of bloating can be broadly classified into lifestyle aspects and medical conditions.

Gynecological Issues

Some women can experience abdominal pain during menstruation. Pregnant women may experience a jabbing discomfort in event of an ectopic pregnancy. This is an issue of pregnancy in which the fertilized egg gets implanted outside the uterus. It could be implanted in the abdomen, ovary or the cervix. This might result in serious pain in the lower right or lower left abdominal area. Inflammation in the fallopian tubes or development of uncomfortable ovarian cysts might also trigger a piercing pain.

“



***The Secret of Easily Preventing Diverticulitis & Achieving Colon Health** Diverticulitis seems to be the new Disease of the Year , in the media. Everywhere you turn you re hearing more doctors and medical studies say More people in the world have either diverticulitis or diverticulosis than ever before! But how many of...*

There is No Rule When It Concerns Normal or Regular Bowel Movements

The general and accepted variety of the frequency of bowel movements is 3 times a day to 3 times a week. If the number is less than three weekly, it is an indicator of constipation. At the exact same time, more than 3 watery stools show diarrhea. There is a possibility, that a person may have one bowel movement a day for a variety of years then the number might increase to two or 3 a day for no evident factor. This is likewise considered to be normal, but would certainly suggest that the frequency of bowel movements for the individual has undergone a change.



[Ibs](#)[Constipation](#)[Irritable Bowel Syndrome](#)[Nausea](#)[Diarrhea](#)[Ibs](#)

Puffed Up Abdomen Causes

An augmentation in the abdominal area can be due to a number of factors, some of which may be because of small health problems, while some may need timely medical interest. Mentioned listed below are a few of the causative factors for stomach bloating. Signs of Abdominal Bloating.

Complications in Pregnancy and Breastfeeding

Consuming aloe vera juice during pregnancy, or while breast feeding is severely limited due to its irritant and purgative properties. Aloe vera ought to be restricted during lactation, as the presence of anthraquinones in the breast milk may cause diarrhea. In pregnant females, aloe vera juice can trigger uterine contractions, as well as result in a spontaneous miscarriage. Aloe vera juice is likewise hazardous for kids below twelve years of age, as it can lead to diarrhea and stomach cramps.

Although there are no magic foods to make your stomach feel better once your symptoms appear, you may be able to decrease the frequency of your symptoms if you follow these pointers:

Eat a Healthy Diet Plan of Low-Fat Foods.

Drink 6 to eight glasses of water daily.

Eat More Regular, Smaller Sized Meals.

Gradually add fiber to your diet (either by taking supplements or consuming high-fiber foods). Soluble fiber, which is in foods like spinach, apples, and figs, can assist with both irregularity and diarrhea. Foods abundant in insoluble fibers, consisting of whole grain breads and cereals, can assist bulk up your stool if you have diarrhea. If you experience gas, you might have added too much fiber too rapidly.

Disclaimer: Content in this document is an advertisement, therefore it may be biased and should not be considered an objective or independent review. Owner of this document may be compensated when you purchase a product by clicking a link in this document. The views and opinions expressed in this document are purely of the author. Any product claim or other representation about a product or service should be verified with the manufacturer, provider or party in question. Any health information in this document is for educational purposes only and is not intended to replace the advice of your health care provider.