

# Pain in Lower Left Abdomen

Abdominal pain can either be a symptom of some disorder or a transient condition in itself. There are some essential organs situated near the abdomen. For that reason, a disorder that happens in any of these organs can result in pain in the abdomen. The pain can be of varying types, persistent or acute, depending on the cause.



## How Supplements can Help

Supplements that consist of probiotics can be an addition to a special diet, or an alternative treatment. \* You can discover probiotic supplements that are specifically designed to help stabilize your digestion health, which will alleviate a great deal of the symptoms that you are experiencing simply by providing a solution to the cause of the concern in the first location. \* Supplements are something that you need to talk to your doctor about and you should always be looking for the most credible probiotics so that you can get the digestive assistance that you deserve. \*

## Ulcer.

Medicine, ulcers are defined as an area of tissue erosion. The causative agents are smoking, stress, and bacterial infection. They may occur in the lining of the gastrointestinal tract. If the left side is affected, then it may lead to mild to severe left side abdominal pain. The region with the ulcer is typically concave in shape due to the fact that of tissue erosion.

## Irritable Bowel Syndrome.

Is often observed that people with irritable bowel syndrome are prone to establishing a bloated stomach. Perhaps, it is one of the very disturbing reasons for bloating. In IBS, there is no problem in particular with the bowel movement, but it's the gut that does not function correctly. For majority, the bloating tends to happen, even get worse, during the evening hours time when one might want to go to social gatherings. Cereal fibers, such as whole bread, oats, and muesli, if avoided, may control the bloating, and significantly reduce the instances of the very same.

- Here are the primary signs present in a person affected by Irritable Bowel Syndrome.
- It changes from every person and may have several of the symptoms but not all of the symptoms may be present.

Apart from the above mentioned ones, other causes consist of disorders like gastroenteritis, abdominal aortic aneurysm, bladder infection, menstrual cramps, ovarian cysts, food poisoning, pelvic inflammatory disease, bowel cancer, transverse colon volvulus, etc. Lower stomach pain in men can be a symptom of prostate conditions, seminal vesiculitis or urinary tract infection. It should be kept in mind that sexually transmitted illness can likewise cause abdominal pain.

## Diverticulitis.

Diverticulitis is a disease of the large intestine. Tenderness and pain in the lower abdomen are the most common symptoms of diverticulitis. This disease is usually observed in older people, above the age of 60.

## Symptoms.

This condition has a number of symptoms. Some suffer from insomnia and loss of appetite. Breakdown makes the person emotionally-drained. This may lead to anxiety attacks or depression. Affected individuals may often show signs of delirium or amnesia. They may have frequent nightmares. Other signs include rapid heartbeat, lightheadedness, irritability, insomnia, shortness of breath, etc. The person experiences fear and stress and anxiety. These signs are covered in the table to offer you a much better idea:

## Ectopic Pregnancy.

Pain in the lower left abdomen during pregnancy can be a popular sign of an ectopic pregnancy. Apart from ectopic pregnancy, it can also be caused due to other factors like ligament pain due to the expanding uterus, luteal cysts, etc. Although abdominal pain is not always related to ectopic pregnancy, seek advice from the doctor if this symptom is observed throughout pregnancy.

- Treatment: Medicines like Methotrexate can help in dealing with early cases of ectopic pregnancy.
- Whereas, if identified in later stages, a surgery may be recommended.

## Endometriosis.

Endometriosis is a disorder observed only in women. In this condition, the tissue (endometrium) that lines the uterus grows outside the uterine cavity. This leads to pain, irritation, advancement of adhesions, etc. Pain during menstruation and urination, including signs like diarrhea, constipation, etc., are observed.

Treatment: In small cases (where discomfort is milder, and no complications are observed), medications, antibiotics, and following a proper diet can be helpful. In case any problems like bowel obstruction are observed, the doctor will recommend hospitalization and surgery.

## You can Naturally Command Your Hypo

"On Dr Litten's system I had no side effects and mostly lots of energy, more energy than I ever had in my life. Getting on the scale every morning is a joy. I have actually lost 28 lbs and Im still losing. Its a pleasure now to buy clothes in small sizes instead of 15-16 or bigger. It's the only diet I have ever been on that I have never felt hungry irritable or tired. My whole life and outlook has changed a new fob, new clothing. I take much much better care of myself than ever before", Janice Mount. Mt Holly New Jersey Dr Litten's system has given me the best results I have ever had out of all the diets, grapefruit diet pills, and Stillman diets.

- Treatment: Smaller stones pass on their own.
- The doctor will advise you to drink fluids, or prescribe medications.
- On the other hand, a surgery helps to get rid of larger kidney stones.

## Ulcerative Colitis.

Crohn's illness and ulcerative colitis are quite similar to each other. Ulcerative colitis is marked by swelling, ulcers, and sores in the inner lining of the colon and rectum. It is a type of inflammatory bowel disease (IBD). Rectal bleeding, abdominal cramping and pain, weight loss, etc., are the symptoms of this condition.

*“ Treatment: This is a chronic disorder, and hence, the symptoms appear gradually. Treatment options include anti-inflammatory medicines, antibiotics, painkillers, etc. Severe cases are treated with surgery. This condition is treated like an autoimmune disorder, and hence, immune system suppressors may even be prescribed.*

Treatment: Making changes in your diet, drinking adequate amounts of water, using laxatives, etc., can help in treating **constipation** in minor cases. In severe cases, if bowel obstruction is diagnosed, surgery may be needed.

- The nature and characteristics of discomfort aid in identifying the exact cause of the pain.
- For that reason, the patient should clearly explain to the doctor whether the pain is dull and aching, or sharp and persistent, acute or chronic.
- A physical examination, X-ray, case history, etc., will be helpful in identifying the underlying causes.

## Eating Behaviors.

Bad eating habits can also lead to stand bloating. So, it is very needed to follow the right eating habits. It includes eating gradually, chewing well, etc. These days due to busy schedule everyone prevents spending time on eating; nevertheless, eating fast makes one swallow extra air that leads to bloating. When one chews food and then swallows it, it gets mixed well with the saliva. This then starts the breakdown of food even before it reaches the digestive tract, thus, helping in digestion.

**Disclaimer: Content in this document is an advertisement**, therefore it may be biased and should not be considered an objective or independent review. Owner of this document may be compensated when you purchase a product by clicking a link in this document. The views and opinions expressed in this document are purely of the author. Any product claim or other representation about a product or service should be verified with the manufacturer, provider or party in question. Any health information in this document is for educational purposes only and is not intended to replace the advice of your health care provider.